

# **WIMPs at Muon colliders**


S.Bottaro, D.Buttazzo, M.Costa, R.Franceschini, P.Panci,  
D.Redigolo, L.Vittorio

2107.09688

Based on: Di Luzio et al. 1810.10993, Han et al. 2009.11287, Capdevilla et al. 2102.11292,  
Bottaro et al. 2103.12766

# Which WIMP?



- **EW** interaction sets abundance: **M** only parameter



# Which WIMP?

- **EW** interaction sets abundance: **M** only parameter
- DM **neutral component** of generic SU(2) n-plet:

$$T_3 + Y = 0$$





# Which WIMP?

- EW interaction sets abundance: **M** only parameter
- DM **neutral component** of generic SU(2) n-plet:

$$T_3 + Y = 0$$

- DD bounds **no Z coupling**:
  - odd n: **Y=0 (real)**
  - even n, **Y≠ 0**: needs mixing partner





# Which WIMP?

- EW interaction sets abundance: **M** only parameter
- DM **neutral component** of generic SU(2) n-plet:

$$T_3 + Y = 0$$

- DD bounds **no Z coupling**:
  - odd n: **Y=0 (real)**
  - even n, **Y ≠ 0**: needs mixing partner
- Real odd-plets **SPECIAL**: IR physics **independent** on completion





# Which WIMP?

- EW interaction sets abundance: **M** only parameter
- DM **neutral component** of generic SU(2) n-plet:

$$T_3 + Y = 0$$

- DD bounds **no Z coupling**:
  - odd n: **Y=0 (real)**
  - even n, **Y ≠ 0**: needs mixing partner
- Real odd-plets **SPECIAL**: IR physics **independent** on completion
- **Computability**: SU(2) Landau Pole  $\gg M \Rightarrow n \leq 13$



# Computing the DM Relic Abundance

Boltzmann equation:

$$\frac{dY}{dx} = -\frac{s(x)}{xH(x)} \langle \sigma v \rangle \left( 1 - \frac{x}{3g_*(x)} \frac{dg_*}{dx} \right) (Y^2(x) - Y_{eq}^2(x))$$

# Computing the DM Relic Abundance

Boltzmann equation:

$$\frac{dY}{dx} = -\frac{s(x)}{xH(x)} \langle \sigma v \rangle \left( 1 - \frac{x}{3g_*(x)} \frac{dg_*}{dx} \right) (Y^2(x) - Y_{eq}^2(x))$$

WHICH CROSS-SECTION?

# Computing the DM Relic Abundance

Boltzmann equation:

$$\frac{dY}{dx} = -\frac{s(x)}{xH(x)} \langle\sigma v\rangle \left(1 - \frac{x}{3g_*(x)} \frac{dg_*}{dx}\right) (Y^2(x) - Y_{eq}^2(x))$$

**WHICH CROSS-SECTION?**

Std. tree-level cross-section:

$$\langle\sigma v\rangle_0 = \frac{\pi\alpha_2^2(2n^4 + 17n^2 - 19)}{16g_\chi M_\chi^2}$$
Correct...

# Computing the DM Relic Abundance

Boltzmann equation:  $\frac{dY}{dx} = -\frac{s(x)}{xH(x)} \langle\sigma v\rangle \left(1 - \frac{x}{3g_*(x)} \frac{dg_*}{dx}\right) (Y^2(x) - Y_{eq}^2(x))$

WHICH CROSS-SECTION?

Std. tree-level cross-section:  $\langle\sigma v\rangle_0 = \frac{\pi\alpha_s^2(2n^4 + 17n^2 - 19)}{16g_\chi M_\chi^2}$       Correct...

... but inaccurate! Important physics is missing

- Sommerfeld enhancement
- Bound states formation

# Computing the DM Relic Abundance

Boltzmann equation:  $\frac{dY}{dx} = -\frac{s(x)}{xH(x)} \langle\sigma v\rangle \left(1 - \frac{x}{3g_*(x)} \frac{dg_*}{dx}\right) (Y^2(x) - Y_{eq}^2(x))$

WHICH CROSS-SECTION?

Std. tree-level cross-section:  $\langle\sigma v\rangle_0 = \frac{\pi\alpha_2^2(2n^4 + 17n^2 - 19)}{16g_\chi M_\chi^2}$       Correct...

... but inaccurate! Important physics is missing

- Sommerfeld enhancement
- Bound states formation

}

Large *non-perturbative, non-relativistic* effects!

# Computing the DM Relic Abundance

Boltzmann equation:  $\frac{dY}{dx} = -\frac{s(x)}{xH(x)} \langle\sigma v\rangle \left(1 - \frac{x}{3g_*(x)} \frac{dg_*}{dx}\right) (Y^2(x) - Y_{eq}^2(x))$

WHICH CROSS-SECTION?

Std. tree-level cross-section:  $\langle\sigma v\rangle_0 = \frac{\pi\alpha_s^2(2n^4 + 17n^2 - 19)}{16g_\chi M_\chi^2}$       Correct...


... but inaccurate! Important physics is missing

- Sommerfeld enhancement
- Bound states formation

Large *non-perturbative, non-relativistic* effects!


keep in mind for later!

# Thermal target results



DM spin	EW n-plet	$M_\chi$ (TeV)
Real scalar	3	$2.53 \pm 0.01$
	5	$15.4 \pm 0.7$
	7	$54.2 \pm 3.1$
	9	$117.8 \pm 15.4$
	11	$199 \pm 42$
	13	$338 \pm 102$
Majorana fermion	3	$2.86 \pm 0.01$
	5	$13.6 \pm 0.8$
	7	$48.8 \pm 3.3$
	9	$113 \pm 15$
	11	$202 \pm 43$
	13	$324.6 \pm 94$


# Thermal target results



DM spin	EW n-plet	$M_\chi$ (TeV)
Real scalar	3	$2.53 \pm 0.01$
	5	$15.4 \pm 0.7$
	7	$54.2 \pm 3.1$
	9	$117.8 \pm 15.4$
	11	$199 \pm 42$
	13	$338 \pm 102$
Majorana fermion	3	$2.86 \pm 0.01$
	5	$13.6 \pm 0.8$
	7	$48.8 \pm 3.3$
	9	$113 \pm 15$
	11	$202 \pm 43$
	13	$324.6 \pm 94$

30-ish TeV collider might probe them

# Thermal target results




DM spin	EW n-plet	$M_\chi$ (TeV)
Real scalar	3	$2.53 \pm 0.01$
	5	$15.4 \pm 0.7$
	7	$54.2 \pm 3.1$
	9	$117.8 \pm 15.4$
	11	$199 \pm 42$
	13	$338 \pm 102$
Majorana fermion	3	$2.86 \pm 0.01$
	5	$13.6 \pm 0.8$
	7	$48.8 \pm 3.3$
	9	$113 \pm 15$
	11	$202 \pm 43$
	13	$324.6 \pm 94$

Need at least 100 TeV collider!

# WIMPs @ high energy lepton colliders

Why high energy lepton colliders?


- **EW nature of signal** : high energy lepton collider as  
**“EW bosons collider”**



# WIMPs @ high energy lepton colliders

Why high energy lepton colliders?


- **EW nature of signal** : high energy lepton collider as  
**“EW bosons collider”**
- **Clean environment**: full event reconstruction  
(**Missing mass**,...)



# WIMPs @ high energy lepton colliders

Why high energy lepton colliders?

- **EW nature of signal** : high energy lepton collider as  
**“EW bosons collider”**
- **Clean environment**: full event reconstruction  
(**Missing mass**,...)




- **More energy in hard cross section**: needed since  
WIMPs are heavy

# How to detect WIMPs @ Muon Collider?

- **Recoils** against invisible objects: **Mono-X,Di-X**

(mono $\gamma$ , monoZ, monoW, mono $\mu$ , Di $\gamma$ , DiW, Di $\mu$ )

Han et al. 2009.11287




Bottaro, MC et al.  
2107.09688

# How to detect WIMPs @ Muon Collider?

- **Recoils** against invisible objects: **Mono-X,Di-X**

(mono $\gamma$ , monoZ, monoW, mono $\mu$ , Di $\gamma$ , DiW, Di $\mu$ )

Han et al. 2009.11287




Bottaro, MC et al.  
2107.09688

- **Disappearing tracks**: lifetime robust prediction

Capdevilla et al.  
2102.11292


Bottaro, MC et al.  
2107.09688



# How to detect WIMPs @ Muon Collider?

- **Recoils** against invisible objects: **Mono-X,Di-X**

(mono $\gamma$ , monoZ, monoW, mono $\mu$ , Di $\gamma$ , DiW, Di $\mu$ )




Han et al. 2009.11287

Bottaro, MC et al.  
2107.09688

- **Disappearing tracks**: lifetime robust prediction


Capdevilla et al.  
2102.11292

Bottaro, MC et al.  
2107.09688




- **Resonances**: Bound States

Bottaro et al.  
2103.12766



# Missing Mass search example: mono- $\gamma$





$$\mu^- \mu^+ \rightarrow \gamma \chi^n \chi^{-n}$$

$$\mu^- \mu^+ \rightarrow \gamma \nu \bar{\nu}$$



# Missing Mass search example: mono- $\gamma$

Han et al.2009.11287



# Missing Mass search example: mono- $\gamma$

Han et al.2009.11287





**S/B<0.1% LOW!**

$$\frac{S}{\sqrt{S + B + (\epsilon_S S)^2 + (\epsilon_B B)^2}}$$

# Missing Mass search example: mono- $\gamma$

Han et al.2009.11287




**S/B<0.1% LOW!**

$\text{MIM} > 2m_\chi$  ,     $|\eta_\gamma| < \eta_{\gamma\text{cut}}$  ,     $\text{MET} > \text{MET}_{\text{cut}}$


$$\frac{S}{\sqrt{S + B + (\epsilon_S S)^2 + (\epsilon_B B)^2}}$$

# Mono-V

Sig





**W**



# Lumi vs Energy (Mono-W)


$$\mathcal{L} \simeq 10 \text{ ab}^{-1} \cdot \left( \frac{\sqrt{s}}{10 \text{ TeV}} \right)^2$$



# Di-V


Sig

Di- $\gamma$



Bkg

Di-W (same sign)





W+W- $\nu\nu$  + mistag

$\epsilon$  mistag=0.1%

Exploit high EW charge of signal

# Mono-X & Di-X results

$\epsilon = 0\%, 0.1\%, 1\%$




**Mono-X:** Mono-W best; S/B<0.1%: need strong pT cuts (E/4-ish)  
**Di-X:** Good for 5; S/B up to 1; robust to  $\epsilon$


# Muon channels

Han et al.  
2009.11287

Sig





Bkg



VBF channels

# Mono- $\mu$ results


Han et al.  
2009.11287



Good at low mass

# Disappearing Tracks

Sig




**Disappearing condition:**  
decay between 5 cm and 12.7 cm

Bkg

BIB hits reconstructed as tracks +  $\gamma$

# Disappearing Tracks

Sig



Bkg

BIB hits reconstructed as tracks +  $\gamma$


**Disappearing condition:**  
decay between 5 cm and 12.7 cm

Recast of Capdevilla et.al  
2102.11292

For details of analysis:  
see next talk by J.Zurita

# Disappearing Tracks

Sig



**Disappearing condition:**  
decay between 5 cm and 12.7 cm

Recast of Capdevilla et.al  
2102.11292

Bkg



BIB hits reconstructed as tracks +  $\gamma$

Splitting/lifetime **FIXED** for real WIMPs

$$M_Q - M_0 \simeq \frac{Q^2 \alpha_{\text{em}} m_W}{2(1 + \cos \theta_W)} : c\tau_{\chi^+} \simeq \frac{120 \text{ mm}}{T(T + 1)}$$

:  
:

# Lumi vs Energy (DT)





**2 $\sigma$ : 35 TeV...**

# Mono-X & Di-X results


$\epsilon = 0\%, 0.1\%, 1\%$

Bottaro, MC et al. 2107.09688c




**1DT:** Good for 3; For 5 comparable to MIM

# Scalar WIMPs



**3S: DTs only hope**

**3S (Disappearing Tracks)**



# Bound States (5F only)

Bottaro et al. 2103.12766

**Peaks in ee cross section**


For indirect searches (virtual corrections) in  $\mu\mu \rightarrow ee$ :  
Di Luzio et al. 1810.10993

# Bound States (5F only)

Bottaro et al. 2103.12766

Peaks in ee cross section

same QN as W's!  
s-channel mixing




# Bound States (5F only)

Bottaro et al. 2103.12766

Peaks in ee cross section

same QN as W's!  
s-channel mixing




$$\sigma(i_1 i_2 \rightarrow B \rightarrow f) \approx \text{BW}(s) \sigma_{\text{peak}}$$

Convoluted with Gaussian beam  
Energy spread  
 $O(1\text{-}10 \text{ GeV})$  vs  $O(0.1\text{-}1 \text{ GeV})$  widths

# Bound States (5F only)


Bottaro et al. 2103.12766



**Discovery in 1 day of running!**

Peaks in ee cross section

same QN as W's!  
s-channel mixing




$$\sigma(i_1 i_2 \rightarrow B \rightarrow f) \approx \text{BW}(s) \sigma_{\text{peak}}$$

Convolved with Gaussian beam  
Energy spread  
 $O(1\text{-}10 \text{ GeV})$  vs  $O(0.1\text{-}1 \text{ GeV})$  widths

# Bound States (5F only)


Bottaro et al. 2103.12766



**Discovery in 1 day of running!**

Peaks in ee cross section


same QN as W's!  
s-channel mixing




$$\sigma(i_1 i_2 \rightarrow B \rightarrow f) \approx \text{BW}(s) \sigma_{\text{peak}}$$

Convolved with Gaussian beam  
Energy spread  
 $O(1\text{-}10 \text{ GeV})$  vs  $O(0.1\text{-}1 \text{ GeV})$  widths

# Collider vs ID/DD



recast from Panci et al.  
1608.00786



# Conclusions

- We computed thermal masses (only parameter) for **real EW WIMPs**

# Conclusions

- We computed thermal masses (only parameter) for **real EW WIMPs**
- Only **3S, 3F, 5F** kinematically allowed if  $\sqrt{s} < 30$  TeV

# Conclusions

- We computed thermal masses (only parameter) for **real EW WIMPs**
- Only **3S, 3F, 5F** kinematically allowed if  $\sqrt{s} < 30$  TeV
- **Mono-X: low S/B**, need high pT cuts. **Mono-W 3F**  $2\sigma$  @ **12 TeV**

# Conclusions

- We computed thermal masses (only parameter) for **real EW WIMPs**
- Only **3S, 3F, 5F** kinematically allowed if  $\sqrt{s} < 30$  TeV
- **Mono-X: low S/B**, need high pT cuts. **Mono-W 3F  $2\sigma$  @ 12 TeV**
- **Di-X**: relevant only for **5plets**. **S/B up to 1**

# Conclusions

- We computed thermal masses (only parameter) for **real EW WIMPs**
- Only **3S, 3F, 5F** kinematically allowed if  $\sqrt{s} < 30$  TeV
- **Mono-X**: **low S/B**, need high pT cuts. **Mono-W 3F**  $2\sigma$  @ **12 TeV**
- **Di-X**: relevant only for **5plets**. **S/B up to 1**
- **Muonic channels**: help for **low mass (3-plets)**

# Conclusions

- We computed thermal masses (only parameter) for **real EW WIMPs**
- Only **3S, 3F, 5F** kinematically allowed if  $\sqrt{s} < 30$  TeV
- **Mono-X**: **low S/B**, need high pT cuts. **Mono-W 3F  $2\sigma$  @ 12 TeV**
- **Di-X**: relevant only for **5plets**. **S/B up to 1**
- **Muonic channels**: help for **low mass (3-plets)**
- **DT**: very good for **3plets**: **3F  $2\sigma$  @ 6 TeV, 3S @ 8 TeV** (only chance)

# Conclusions

- We computed thermal masses (only parameter) for **real EW WIMPs**
- Only **3S, 3F, 5F** kinematically allowed if  $\sqrt{s} < 30$  TeV
- **Mono-X**: **low S/B**, need high pT cuts. **Mono-W 3F  $2\sigma$  @ 12 TeV**
- **Di-X**: relevant only for **5plets**. **S/B up to 1**
- **Muonic channels**: help for **low mass (3-plets)**
- **DT**: very good for **3plets**: **3F  $2\sigma$  @ 6 TeV, 3S @ 8 TeV** (only chance)
- **5F**: **resonance** of  $n=1$  BS. **Discovery** in 1 day provided 0.01% energy resol



# Thanks for the attention!

# Backup

# Mono-X

$\epsilon=0\%$



Bottaro, MC et al. 2107.09688




# Di-X

$\epsilon=0\%$


Bottaro, MC et al. 2107.09688c




# Disappearing Tracks



# Scalar WIMPs



## 5S (Disappearing Tracks)




$2\sigma$ : hopeless

# Cuts


	$\sqrt{s}$	$\epsilon_{\text{sys}}$	Majorana 3-plet					Majorana 5-plet				
			$\eta_X^{\text{cut}}$	$p_{T,X}^{\text{cut}}$ [TeV]	$S_{95\%}$	$S_{95\%}/B$	$M_{95\%}$ [TeV]	$\eta_X^{\text{cut}}$	$p_{T,X}^{\text{cut}}$ [TeV]	$S_{95\%}$	$S_{95\%}/B$	$M_{95\%}$ [TeV]
Mono- $\gamma$	3 TeV	0	2.4	0.18	1007	0.004	0.72	2.4	0.0	3038	0.001	1.4
		1%	2.2	0.24	746	0.006	0.67	1.2	0.0	3683	0.003	1.3
		1%	1.2	0.78	107	0.05	0.58	0.6	0.3	639	0.02	1.1
	14 TeV	0	1.6	2.5	360	0.01	2.2	2.2	0.28	3693	0.001	5.5
		1%	1.6	2.8	323	0.01	2.2	1.2	0.84	1300	0.004	5.2
		1%	1.0	4.5	108	0.05	1.9	0.8	2.8	331	0.03	4.4
	30 TeV	0	1.2	7.8	174	0.02	4.4	1.6	1.8	1795	0.002	11
		1%	1.2	7.8	175	0.02	4.4	1.0	2.4	1312	0.004	11
		1%	1.2	8.4	190	0.03	4.0	0.8	6.0	455	0.03	8.8
Mono-W (inclusive)	3 TeV	0	1.6	0.36	842	0.005	0.79	2.2	0.06	5625	0.0007	1.2
		1%	1.4	0.48	534	0.008	0.78	1.0	0.24	1649	0.004	1.2
		1%	1.0	0.84	172	0.04	0.64	0.6	0.54	515	0.02	1.0
	14 TeV	0	1.6	2.0	819	0.005	3.4	1.8	0.56	5325	0.0008	5.5
		1%	1.6	2.2	665	0.007	3.3	1.0	1.4	1342	0.004	5.2
		1%	0.8	4.2	155	0.04	2.8	1.2	2.5	635	0.03	4.4
	30 TeV	0	1.4	5.4	696	0.006	6.7	1.8	1.8	3946	0.001	12
		1%	1.4	5.4	606	0.007	6.7	1.4	2.4	2771	0.003	11
		1%	1.0	9.0	211	0.03	5.2	0.8	5.4	813	0.02	9.3

# FCC-hh prospects

Han et al.  
1805.00015







mono-jet



DT

# Indirect collider prospects

$\mu\mu \rightarrow ff$   
corrections




Expected 95% CL exclusion limits

Di Luzio et al.  
1810.10993

# Integrated luminosity for BS

Bottaro et al. 2103.12766



# DT recast

$$P(\theta, r_{\min}, r_{\max}) = \int_{r_{\min}}^{r_{\max}} \frac{dr}{c\tau\beta\gamma \sin\theta} \epsilon_{\text{rec}}(r, \theta) e^{-r/(c\tau\beta\gamma \sin\theta)},$$

Capdevilla et al.  
2102.11292

Requirement / Region	SR <sub>1t</sub> <sup>γ</sup>	SR <sub>2t</sub> <sup>γ</sup>
Veto	leptons and jets	
Leading tracklet $p_T$ [GeV]	> 300	> 20
Leading tracklet $\theta$ [rad]	[2/9π, 7/9π]	
Subleading tracklet $p_T$ [GeV]	-	> 10
Tracklet pair $\Delta z$ [mm]	-	< 0.1
Photon energy [GeV]	> 25	> 25